

Compatibility of $\alpha_s(M_\tau)$ with $\alpha_s(M_Z)$

T.G. STEELE

*Department of Physics and Engineering Physics and
Saskatchewan Accelerator Laboratory
University of Saskatchewan
Saskatoon, Saskatchewan S7N 5C6, Canada.*

V. ELIAS

*Department of Applied Mathematics, The University of Western Ontario,
London, Ontario N6A 5B7, Canada.*

February 1, 2008

Abstract

The current phenomenological determinations of $\alpha_s(M_\tau)$ and $\alpha_s(M_Z)$ are shown to be only marginally consistent with the QCD evolution of the strong coupling constant between M_Z and M_τ . This motivates a revised estimate of $\alpha_s(M_\tau)$ since the perturbative series used to extract $\alpha_s(M_\tau)$ from the τ hadronic width exhibits slow convergence. Padé summation techniques provide an estimate of these unknown higher-order effects, leading to the revised determination $\alpha_s(M_\tau) = 0.333 \pm 0.030$. This value is 10% smaller than current estimates, improving the compatibility of phenomenological estimates for $\alpha_s(M_\tau)$ and $\alpha_s(M_Z)$ with the QCD evolution of the strong coupling constant.

The Particle Data Group (PDG) quotes the following values for the strong coupling constant as determined from Z^0 and τ decays [1].

$$\alpha_s(M_\tau) = 0.370 \pm 0.033 \tag{1}$$

$$\alpha_s(M_Z) = 0.118 \pm 0.003 \tag{2}$$

Since these determinations of α_s occur at such widely separated energies, the compatibility of these values of α_s with the QCD evolution of the coupling constant is an important test of both QCD and the phenomenological results used to extract the coupling constant from the experimental data. In particular, $\alpha_s(M_\tau)$ is sufficiently large that presently unknown terms from higher order perturbation theory could substantially alter the value of $\alpha_s(M_\tau)$ extracted from the experimental data. Padé approximant methods provide estimates of the aggregate effect of (presently unknown) terms from higher-order perturbation theory [2, 3, 4, 5]. As shown below, the use of Padé summation to estimate such terms leads to a substantial decrease in the value of $\alpha_s(M_\tau)$ extracted from τ decays, improving the compatibility of $\alpha_s(M_\tau)$ and $\alpha_s(M_Z)$ with the QCD evolution of the coupling constant.

The QCD evolution of the coupling constant is governed by the β function which is now known to 4-loop order [6]. Using the conventions of [7], $a \equiv \frac{\alpha_s}{\pi}$ satisfies the differential equation

$$\mu^2 \frac{da}{d\mu^2} = \beta(a) = -a^2 \sum_{i=0}^{\infty} \beta_i a^i \quad , \quad a \equiv \frac{\alpha_s}{\pi} \tag{3}$$

$$\beta_0 = \frac{11 - \frac{2}{3}n_f}{4} \quad , \quad \beta_1 = \frac{102 - \frac{38}{3}n_f}{16} \quad , \quad \beta_2 = \frac{\frac{2857}{2} - \frac{5033}{18}n_f + \frac{325}{54}n_f^2}{64} \quad (4)$$

$$\beta_3 = 114.23033 - 27.133944n_f + 1.5823791n_f^2 - 5.8566958 \times 10^{-3}n_f^3 \quad (5)$$

Using the value $\alpha_s(M_Z)$ as an initial condition, the coupling constant can be evolved to the desired energy using the differential equation (3). The only subtlety in this approach is the location of flavour thresholds where the number of effective flavour degrees of freedom n_f change. In general, matching conditions must be imposed at these thresholds to relate QCD with n_f quarks to an effective theory with $n_f - 1$ light quarks and a decoupled heavy quark [8]. Using the matching threshold μ_{th} defined by $m_q(\mu_{th}) = \mu_{th}$, where m_q is the running quark mass, the matching condition to three-loop order is [9]

$$a^{(n_f-1)}(\mu_{th}) = a^{(n_f)}(\mu_{th}) \left[1 + 0.1528 \left[a^{(n_f)}(\mu_{th}) \right]^2 + \{0.9721 - 0.0847(n_f - 1)\} \left[a^{(n_f)}(\mu_{th}) \right]^3 \right] \quad (6)$$

leading to a discontinuity of α_s across the threshold. Thus to determine the coupling constant at energies between the c quark threshold and the b quark threshold, the β function with $n_f = 5$ is used to run $\alpha_s^{(5)}$ from M_Z to $\mu_{th} = m_b(\mu_{th}) \equiv m_b$ using (2) as an initial condition. The matching condition (6) is then imposed to find the value of $\alpha_s^{(4)}(m_b)$ which is then used as an initial condition to evolve α_s to lower energies via the $n_f = 4$ β function.

If $\alpha_s(M_Z)$ is used as the input value to determine the QCD prediction of $\alpha_s(M_\tau)$, then one might legitimately be concerned about the effect of (unknown) higher-order terms in the β function at lower energies where α_s is larger. Padé approximations have proven their utility in determining higher-order terms in the β function. For example, using as input the four-loop β function in $O(N)$ gauge theory [10], asymptotic Padé methods described in Section II of [4] are able to predict the five-loop term to better than 10% of the known five-loop contributions for $N \leq 4$ [3, 11]. When these same methods are applied to QCD, the following predictions for the unknown five-loop contribution to the β function are obtained [11].

$$n_f = 4 : \quad \beta_4 = 83.7563 \quad (7)$$

$$n_f = 5 : \quad \beta_4 = 134.56 \quad (8)$$

From these predictions, β functions containing [2|2] Padé approximants can be constructed to estimate the sum of all higher-order contributions. These Padé-summations, whose Maclaurin expansions reproduce β_1 , β_2 , β_3 and the asymptotic Padé-approximant estimates (7,8) of β_4 , are given by:

$$n_f = 4 : \quad \beta(a) = -\frac{25x^2}{12} \left[\frac{1 - 5.8963a - 4.0110a^2}{1 - 7.4363a + 4.3932a^2} \right] \quad (9)$$

$$n_f = 5 : \quad \beta(a) = -\frac{23x^2}{12} \left[\frac{1 - 5.9761a - 6.9861a^2}{1 - 7.2369a - 0.66390a^2} \right] \quad (10)$$

Thus the QCD prediction of $\alpha_s(M_\tau)$ depends on only two parameters: the initial condition $\alpha_s(M_Z)$ and the position of the five-flavour threshold defined by $m_b(m_b) = m_b$. As will be discussed below, the uncertainty in the Particle Data Group value [1] for this threshold

$$m_b(m_b) = (4.3 \pm 0.3)\text{GeV} \quad (11)$$

has a negligible effect on the QCD prediction of $\alpha_s(M_\tau)$ compared with the uncertainty in $\alpha_s(M_Z)$ (2).

The compatibility of the experimentally/phenomenologically determined values $\alpha_s(M_Z)$ and $\alpha_s(M_\tau)$ with the QCD evolution of the coupling constant can now be studied. Figure 1 shows the effect on $\alpha_s(Q)$ of progressive increases in the number of perturbative terms in the β function, culminating with the Padé summation (9,10) for β . It is evident that the curves for $\alpha_s(Q)$ appear to converge from below to that generated by the Padé summation of the β function, since the gaps between curves of successive order decrease. Using the input values (2,11) for the QCD evolution of α_s down from the Z^0 to τ mass, we obtain the following range of values for $\alpha_s(M_\tau)$ for successive orders of perturbation theory [11]:

$$2 - \text{loop} \quad 0.2910 \leq \alpha_s(M_\tau) \leq 0.3391 \quad (12)$$

$$3 - \text{loop} \quad 0.2944 \leq \alpha_s(M_\tau) \leq 0.3451 \quad (13)$$

$$4 - \text{loop} \quad 0.2957 \leq \alpha_s(M_\tau) \leq 0.3477 \quad (14)$$

$$\text{Padé summation} \quad 0.2963 \leq \alpha_s(M_\tau) \leq 0.3489 \quad (15)$$

The dominant effect on the uncertainty in the above originates from $\alpha_s(M_Z)$ — the effect of the uncertainty in the five-flavour threshold (11) is inconsequential. Thus as illustrated in Figure 2, the empirical determinations of α_s at M_Z and M_τ are only marginally consistent with the QCD evolution of the coupling constant. This motivates a revised determination of $\alpha_s(M_\tau)$ using Padé approximation techniques.

The seminal QCD analysis of the τ hadronic width [12]

$$R_\tau \equiv \frac{\Gamma(\tau \rightarrow \nu_\tau + \text{hadrons})}{\Gamma(\tau \rightarrow \nu_\tau + e + \bar{\nu}_e)} \quad (16)$$

leads to the following result for R_τ .

$$R_\tau = 3 \left(|V_{ud}|^2 + |V_{us}|^2 \right) S_{EW} \left[1 + \delta'_{EW} + \delta^{(0)} + \sum_{D=2,4,6} \cos^2 \theta_c \delta_{ud}^{(D)} + \sin^2 \theta_c \delta_{us}^{(D)} \right] \quad (17)$$

To facilitate comparison of the Padé improvement of this result with previous research we use values for the parameters in (17) identical to [12]: the CKM matrix elements,

$$|V_{ud}| = 0.9753 \quad , \quad |V_{us}| = 0.2210 \quad (18)$$

and the Cabibbo angle

$$\sin^2 \theta_c = \frac{|V_{us}|^2}{|V_{ud}|^2 + |V_{us}|^2} \quad (19)$$

The electroweak contributions in (17) are

$$S_{EW} = 1.0194 \quad , \quad \delta'_{EW} = 0.001 \quad . \quad (20)$$

The mass-independent perturbative contribution to R_τ is the dominant effect: in the $\overline{\text{MS}}$ scheme, it takes the form

$$1 + \delta^{(0)} = 1 + a(M_\tau) + 5.2023 [a(M_\tau)]^2 + 26.366 [a(M_\tau)]^3 \quad . \quad (21)$$

Note that for a characteristic value $\alpha_s(M_\tau) = 0.37$, the contributions of various orders to $\delta^{(0)}$ are

$$1 + \delta^{(0)} = 1 + 0.118 + 0.072 + 0.043 \quad (22)$$

illustrating the slow convergence of the perturbation series, and hence the potentially important role of presently unknown terms from higher-order perturbation theory.

Perturbative quark mass contributions to R_τ occur for $D = 2$ in the summation. The quantity $\delta_{ij}^{(2)}$, representing the average of the axial and vector contributions to R_τ , is

$$\delta_{ij}^{(2)} = -8 \left[1 + \frac{16}{3} a(M_\tau) \right] \frac{m_i^2(M_\tau) + m_j^2(M_\tau)}{M_\tau^2} \quad (23)$$

where $m_i(M_\tau)$ denotes the running quark mass (of flavour i) which has *implicit* dependence on $\alpha_s(M_\tau)$ through its running behaviour

$$m_i(\mu) = \hat{m}_i [2\beta_0 a(\mu)]^{\frac{\gamma_0}{2\beta_0}} \left[1 + \frac{\beta_1}{\beta_0} \left(-\frac{\gamma_0}{2\beta_0} + \frac{\gamma_1}{2\beta_1} \right) a(\mu) \right] \quad (24)$$

$$\gamma_0 = 2 \quad , \quad \gamma_1 = \frac{101}{12} - \frac{5}{18} n_f \quad (25)$$

To facilitate comparison of our results with [12], we use identical values for the RG invariant quark masses $\hat{m}_i$.

$$\hat{m}_u = 8.7 \text{ MeV} \quad , \quad \hat{m}_d = 15.4 \text{ MeV} \quad , \quad \hat{m}_s = 270 \text{ MeV} \quad (26)$$

The total effect of these $D = 2$ mass corrections is small in comparison with $\delta^{(0)}$ since the potentially large s quark contributions are suppressed by $\sin \theta_c$ in (17).

Dimension-four ($D = 4$) and higher contributions to R_τ are dominantly non-perturbative in origin, and involve the QCD condensates which originate from the operator-product expansion [13]. The average of the axial and vector contributions for $D = 4$ is

$$\begin{aligned} \delta_{ij}^{(4)} &= \frac{11\pi^2 a^2(M_\tau)}{4M_\tau^4} \langle aGG \rangle + \frac{16\pi^2}{M_\tau^4} \left(1 + \frac{27}{8} a^2(M_\tau) \right) \langle m_i \bar{\psi}_i \psi_i + m_j \bar{\psi}_j \psi_j \rangle \\ &\quad - \frac{8\pi^2 a^2(M_\tau)}{M_\tau^4} \sum_{k=u,d,s} \langle m_k \bar{\psi}_k \psi_k \rangle + \frac{24m_i^2(M_\tau) m_j^2(M_\tau)}{M_\tau^4} \\ &\quad + \left(-\frac{48}{7a(M_\tau)} + \frac{22}{7} \right) \frac{m_i^4(M_\tau) + m_j^4(M_\tau)}{M_\tau^4} \end{aligned} \quad (27)$$

which has explicit and implicit dependence on $\alpha_s(M_\tau)$. Values for the QCD condensates appearing in (27), as used in [12], are

$$\langle aGG \rangle = 0.02 \text{ GeV}^4 \quad (28)$$

$$\langle m_j \bar{\psi}_j \psi_j \rangle \equiv -\hat{m}_j \hat{\mu}_j^3 \quad , \quad \hat{\mu}_d = \hat{\mu}_u = 189 \text{ MeV} \quad , \quad \hat{\mu}_s = 160 \text{ MeV} \quad (29)$$

Dimension-six contributions are the dominant non-perturbative effect. After use of the effective phenomenological equivalent of the vacuum saturation hypothesis [13] parametrized by the (near) RG-invariant quantity,

$$\rho \alpha_s \langle \bar{\psi} \psi \rangle^2 = 3.8 \times 10^{-4} \text{ GeV}^6 \quad (30)$$

the average of the axial and vector $D = 6$ contributions is [12]

$$\delta_{ij}^{(6)} = -\frac{512\pi^3}{27M_\tau^6} \rho \alpha_s \langle \bar{\psi} \psi \rangle^2 = -0.007 \quad (31)$$

Dimension-eight and higher dimensional non-perturbative contributions to R_τ are negligible [12].

Padé approximation techniques can now be applied to the determination of $\alpha_s(M_\tau)$ from an experimentally determined R_τ . Asymptotic Padé approximation (APAP) techniques use convergence properties of the perturbative expansion to provide improved estimates of the fifth order term in a perturbation series given knowledge of the fourth-order series [4]. Explicit field-theoretical tests of the APAP approach where the fifth order terms are known lead to extremely accurate results. For example, the $N = 1$ limit of the five-loop β function for $O(N)$ gauge theory has been calculated explicitly [10]. Using the APAP algorithm described in Section II of [4], it is possible to make an APAP estimate of the known five-loop term β_5 based upon the contributions from the previous four terms β_1 – β_4 . This estimate has been found to be well within 0.2% of the answer obtained by direct calculation [10].

Applying the same APAP algorithm to the perturbation series (21) leads to the following prediction for the $[a(M_\tau)]^4$ perturbative contribution:

$$1 + \delta^{(0)} = 1 + a(M_\tau) + 5.2023 [a(M_\tau)]^2 + 26.366 [a(M_\tau)]^3 + 132.44 [a(M_\tau)]^4 \quad (32)$$

It is significant to note that this prediction is very close to the *maximum* estimated size of the fourth order effect used to determine the theoretical uncertainty in [12], indicating an underestimate of the higher order effects in previous work. The APAP result in (32) should be compared with that obtained previously from (non-asymptotic) Padé methods [5]

$$1 + \delta^{(0)} = 1 + a(M_\tau) + 5.2023 [a(M_\tau)]^2 + 26.366 [a(M_\tau)]^3 + 109.2 [a(M_\tau)]^4 \quad (33)$$

The coefficient of the a^4 term is 20% smaller in the non-asymptotic prediction.

To determine the effect of the APAP prediction (32) on $\delta^{(0)}$ we again consider a characteristic value $\alpha_s(M_\tau)$

$$\delta^{(0)} = 0.118 + 0.072 + 0.043 + 0.025 \quad (34)$$

The convergence of this series is still quite slow: the $\mathcal{O}(a^4)$ contribution is 20% of the leading perturbative contribution. Such slow convergence indicates that the further higher-order terms could have a significant effect on $\delta^{(0)}$. A Padé summation, in this case a [2|2] Padé approximant, provides a means of estimating the total effect of higher order terms in the perturbation series. Using the APAP determination in (32) leads to the following [2|2] Padé approximation for $\delta^{(0)}$.

$$1 + \delta^{(0)} = \frac{1 - 6.5483a(M_\tau) + 10.5030 [a(M_\tau)]^2}{1 - 7.5483a(M_\tau) + 12.8514 [a(M_\tau)]^2} \quad (35)$$

Effects of the Padé improvement of $\delta^{(0)}$ on the $\alpha_s(M_\tau)$ dependence of R_τ , and hence the determination of $\alpha_s(M_\tau)$, can now be investigated in both the truncated form (32) or [2|2] summation form (35). Figure 3 compares the dependence of R_τ on $\alpha_s(M_\tau)$ using the full $D = 2, 4, 6$ contributions [equations (23,27,31) and electroweak effects (20) for the four different scenarios for $\delta^{(0)}$: the known perturbation series (21), the non-asymptotic Padé expression (33), the asymptotic Padé (APAP) expression (32), and the Padé summation (35). It is evident that at the PDG value $a(M_\tau) = 0.370/\pi = 0.117$, the Padé effects lead to a significant *increase* in R_τ . In comparison with the perturbative result, the size of the R_τ enhancement obtained from the [2|2] Padé summation (35) is roughly twice the enhancement obtained directly from (32,33), indicating the significance of the higher-order effects estimated in the [2|2] Padé summation. To

facilitate comparison with Table 4 in [12] and for future determination of $\alpha_s(M_\tau)$, Table 1 contains R_τ for selected values of $\alpha_s(M_\tau)$ in the four scenarios ¹.

The Padé enhancement of R_τ implies that the value of $\alpha_s(M_\tau)$ extracted from the experimental measurement of R_τ will *decrease*. This is illustrated in Figure 4 which inverts the relation between R_τ and $\alpha_s(M_\tau)$. Again, in comparison with the perturbative result, the decrease in $\alpha_s(M_\tau)$ following from the Padé summation (35) is roughly twice the decrease one obtains from the Padé estimate (32,33) of just the next-order contribution to $\delta^{(0)}$. For the perturbatively-extracted PDG value $\alpha_s(M_\tau) = 0.37$, the Padé summation would lead to a decrease of approximately 10% to $\alpha_s(M_\tau) = 0.33$.

In conclusion, Padé summation methods indicate that the PDG central value $\alpha_s(M_\tau) = 0.37$ should be brought down by approximately 10% to include the estimated effect of higher order QCD corrections. Assuming the same relative uncertainty characterizing the PDG estimate, we find that

$$\alpha_s(M_\tau) = 0.333 \pm 0.030 \tag{36}$$

a result in much closer agreement with the ranges (14,15) devolving from $\alpha_s(M_Z)$ via four-loop and Padé-improved four-loop QCD β functions. Thus, Padé methods are seen to improve the compatibility of $\alpha_s(M_\tau)$ and $\alpha_s(M_Z)$ with the coupling constant evolution anticipated from QCD.

Acknowledgements: The authors are grateful for the financial support of the Natural Sciences and Engineering Research Council of Canada (NSERC).

¹The agreement of the “perturbative” column of the Table with [12] provides a consistency check on our calculations.

References

- [1] Particle data Group, R.M. Barnett *et al* Phys. Rev. **D54** (1996) 1.
- [2] M.A. Samuel, G. Li, E. Steinfelds, Phys. Rev. **D48** (1993) 869; M.A. Samuel, G. Li, E. Steinfelds, Phys. Rev. **E51** (1995) 3911.
- [3] J. Ellis, M. Karliner, M.A. Samuel, Phys. Lett. **B400** (1997) 176.
- [4] J. Ellis, I. Jack, D.R.T. Jones, M. Karliner, M.A. Samuel, Phys. Rev **D57** (1998) 2665.
- [5] M.A. Samuel, J. Ellis, M. Karliner, Phys. Rev. Lett. **74** (1995) 3911.
- [6] T. van Ritbergen, J.A.M. Vermaseren, S.A. Larin, Phys. Lett. **B400** (1997) 379.
- [7] K.G. Chetyrkin, Phys. Lett. **B404** (1997) 161.
- [8] G. Rodrigo, A. Santamaria, Phys. Lett. **B313** (1993) 441.
- [9] K.G. Chetyrkin, B.A. Kniehl, M. Steinhauser, hep-ph/9708255 v2
- [10] H. Kleinart, J. Neu. V. Schulte-Frohlinde, K.G. Chetyrkin, S.A. Larin, Phys. Lett. **B272** (1991) 36, Phys. Lett. **B319** (1993) 545 Erratum.
- [11] V. Elias, T.G. Steele, F. Chistie, R. Migneron, K. Sprague, hep-ph/9806324 .
- [12] E. Braaten, S. Narison, A. Pich, Nucl. Phys. **B373** (1992) 581.
- [13] M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, Nucl. Phys. **B147** (1979) 385.

$\alpha_s(M_\tau)$	R_τ			
	Perturbative	Nonasymptotic Padé	APAP	Padé Summation
.20	3.295	3.300	3.301	3.305
.21	3.314	3.320	3.322	3.326
.22	3.333	3.341	3.343	3.348
.23	3.353	3.362	3.364	3.371
.24	3.374	3.385	3.387	3.396
.25	3.395	3.408	3.411	3.422
.26	3.417	3.433	3.436	3.449
.27	3.440	3.458	3.462	3.478
.28	3.463	3.484	3.489	3.509
.29	3.487	3.511	3.517	3.542
.30	3.512	3.540	3.546	3.576
.31	3.538	3.570	3.576	3.613
.32	3.564	3.600	3.608	3.653
.33	3.592	3.632	3.641	3.695
.34	3.620	3.666	3.675	3.740
.35	3.649	3.700	3.711	3.789
.36	3.678	3.736	3.748	3.841
.37	3.709	3.773	3.787	3.897
.38	3.741	3.812	3.827	3.958
.39	3.773	3.852	3.869	4.024
.40	3.806	3.894	3.913	4.097
.41	3.841	3.937	3.958	4.175
.42	3.876	3.982	4.005	4.262
.43	3.912	4.029	4.054	4.357
.44	3.949	4.077	4.105	4.469

Table 1: Values of R_τ for selected $\alpha_s(M_\tau)$. The columns differ only in the treatment of $\delta^{(0)}$ used in R_τ : the “perturbative” column uses (21), the “Padé summation” column uses (35), the “APAP” column uses (32), and the “Nonasymptotic Padé” uses (33).




Figure 1: Effect of increasing the order of perturbation theory in the QCD evolution of the strong coupling constant using $\alpha_s(M_Z)$ as an initial condition. Higher-loop terms in the β function progressively increase α_s from the 2-loop order bottom (solid) curve to the Padé summation top (dashed-dotted) curve, sandwiching the three- and four-loop curves.




Figure 2: Comparison of $\alpha_s(M_\tau)$ extracted from experiment (right data point) and $\alpha_s(M_\tau)$ determined by QCD evolution of α_s from M_Z to M_τ (left data point).




Figure 3: R_τ as a function of $\alpha_s(M_\tau)$ using the four different treatments of $\delta^{(0)}$. The solid curve uses the perturbative expression (21), the dotted curve uses the non-asymptotic Padé (33), the dashed-dotted curve uses the APAP expression (32), and the dashed curve uses the Padé summation (35).




Figure 4: $\alpha_s(M_\tau)$ as a function of R_τ using the four different treatments of $\delta^{(0)}$. The solid curve uses the perturbative expression (21), the dotted curve uses the non-asymptotic Padé (33), the dashed-dotted curve uses the APAP expression (32), and the dashed curve uses the Padé summation (35).